TANTIA UNIVERSITY JOURNAL OF HOMOEOPATHY AND MEDICAL SCIENCE
Website- www.tjhms.com Email : tujhms@tantiauniversity.com
Peer-Reviewed Journal, E-ISSN: 2581-8899, P-ISSN: 2581-978X
Volume 8|Issue 3|July –Sept. 2025|
REVIEW ARTICLE

BRACHIAL PLEXUS: ANATOMY, CLINICAL CORRELATIONS, AND HOMEOPATHY

Dr Hemlata Singhal
Professor, Department of Anatomy, Homeopathic Medical College

Abstract

Received- 15/09/2025
Revised- 29/09/2025
Accepted– 30/09/2025

Key Word- Brachial plexus, Erb's palsy, Klumpke's palsy, Anatomy

Corresponding Author:-
Dr Hemlata Singhal
Professor, Department of Anatomy, Homeopathic Medical College

The brachial plexus is a vital network of nerves that originates from the spinal cord in the neck and supplies the upper limb. It is responsible for motor and sensory innervation of the shoulder, arm, forearm, and hand. Injuries or pathologies affecting the brachial plexus can lead to significant disability, including weakness, paralysis, or sensory loss. Conventional medical management often includes surgical intervention, and intensive physiotherapy to restore limb function. However, the prognosis can be variable and recovery may be incomplete. In this context, homeopathy offers a complementary therapeutic approach, focusing on individual symptomatology, enhancing the body's innate healing mechanisms, and addressing associated pain, stiffness, or weakness. Remedies such as Hypericum perforatum, Causticum, and Rhus toxicodendron have traditionally been used for conditions involving nerve injuries and neuropathic pain.

INTRODUCTION

The brachial plexus is a highly intricate network of peripheral nerves that arises from the anterior rami of the lower cervical and upper thoracic spinal nerves (C5–T1). It serves as the major conduit for motor and sensory innervation to the upper limb, shoulder girdle, and part of the chest wall. The term "plexus" refers to a branching network of nerves, and the brachial plexus is among the most clinically significant of all peripheral nerve plexuses due to its role in controlling the functional integrity of the upper extremity.

Historically, the anatomical description of the brachial plexus has been credited to early anatomists of the 18th and 19th centuries, who recognized its importance in surgical practice and trauma management. Its anatomical complexity lies not only in its formation through roots, trunks, divisions, cords, and branches, but also in its wide distribution, which ensures fine motor control and sensory precision of the hand—features unique to humans and vital for day-to-day activities.

Clinically, the brachial plexus is highly vulnerable to trauma, obstetric injury, inflammation, neoplastic invasion, and compression syndromes. Conditions such as Erb-Duchenne palsy, Klumpke's palsy can severely compromise limb function, leading to paralysis, sensory deficits, and chronic pain. These disorders may affect individuals across all age groups—from neonates with birth-related injuries to adults sustaining high-velocity trauma. (1-2)

ANATOMY OF THE BRACHIAL PLEXUS

The brachial plexus is a complex network of nerves that originates from the anterior rami of spinal nerves C5 to T1. It is responsible for the motor and sensory innervation of the upper limb, including the shoulder, arm, forearm, and hand. The plexus is anatomically divided into five key components: roots, trunks, divisions, cords, and branches.

1. Roots

- The roots are formed by the anterior rami of spinal nerves C5, C6, C7, C8, and T1.
- These roots emerge from the spinal cord and pass between the anterior and middle scalene muscles in the neck.
- Important nerves arising from the roots include:
  - Dorsal scapular nerve (C5): innervates rhomboid muscles and levator scapulae.
  - Long thoracic nerve (C5–C7): innervates the serratus anterior muscle.

2. Trunks

- The roots converge to form three trunks:
  - Superior trunk: formed by C5 and C6
  - Middle trunk: continuation of C7
  - Inferior trunk: formed by C8 and T1
- These trunks lie in the posterior triangle of the neck.
- Branches from trunks include:
  - Suprascapular nerve: innervates supraspinatus and infraspinatus
  - Nerve to subclavius

3. Divisions

Each trunk splits into the Anterior division and Posterior division. This results in six divisions (three anterior and three posterior), which pass into the axilla.

4. Cords

- The divisions regroup around the axillary artery to form three cords:
  - ➢ Lateral cord: formed by anterior divisions of superior and middle trunks.
  - ➢ Posterior cord: formed by all three posterior divisions.
  - ➢ Medial cord: formed by anterior division of the inferior trunk.
- Each cord gives rise to several branches:
  - ➢ Lateral cord: lateral pectoral nerve, musculocutaneous nerve, lateral root of median nerve.
  - ➢ Medial cord: medial pectoral nerve, ulnar nerve, medial cutaneous nerves of arm and forearm, medial root of median nerve
  - ➢ Posterior cord: axillary nerve, radial nerve, upper and lower subscapular nerves, thoracodorsal nerve

5. Terminal Branches

These are the five major nerves that continue into the upper limb:

1. **Musculocutaneous nerve (C5–C7):**
   - ➢ Motor: biceps brachii, brachialis, coracobrachialis
   - ➢ Sensory: lateral forearm
2. **Axillary nerve (C5–C6):**
   - ➢ Motor: deltoid, teres minor
   - ➢ Sensory: lateral shoulder ("regimental badge" area)
3. **Radial nerve (C5–T1):**
   - ➢ Motor: triceps brachii, forearm extensors
   - ➢ Sensory: posterior arm, forearm, and hand
4. **Median nerve (C6–T1):**
   - ➢ Motor: forearm flexors, thenar muscles, lateral lumbricals
   - ➢ Sensory: lateral palm and fingers
5. **Ulnar nerve (C8–T1):**
   - ➢ Motor: intrinsic hand muscles, flexor carpi ulnaris, part of flexor digitorum profundus
   - ➢ Sensory: medial hand and fingers



Figure 1: Anatomy of Brachial Plexus

Functions Of The Brachial Plexus

1. **Motor Function:** Controls nearly all muscles of the shoulder girdle and upper limb, enabling actions like lifting, grasping, and fine finger movements.

2. **Sensory Function:** Provides sensation to the skin over the shoulder, arm, forearm, and hand.
3. **Autonomic Function:** Supplies sympathetic fibers to blood vessels, sweat glands, and skin of the upper limb. (3-4)

Clinical Relevance

- Erb-Duchenne Palsy (upper trunk injury): caused by traction on neck; affects shoulder and elbow flexors.
- This leads to the classical "waiter's tip" position:
    - Arm hangs by the side (loss of abduction)
    - Arm medially rotated (loss of external rotation)
    - Forearm pronated (loss of supination)
    - Elbow extended (loss of flexion)

Figure 2: Erb's Palsy

- Klumpke's palsy (lower trunk injury): often from forceful arm pulling; affects hand muscles and may cause Horner's syndrome.
- This results in the "claw hand" deformity:
    - Hyperextension of the metacarpophalangeal joints
    - Flexion of the interphalangeal joints (5-6)

Figure 3: Klumpke's Palsy

ROLE OF HOMEOPATHY MEDICINE

Homeopathy aims to stimulate the body's natural healing mechanism and may be supportive in brachial plexus injuries or neuropathies. Remedies are selected based on the totality of symptoms, including type of pain, degree of weakness, and mental state. Some commonly indicated medicines include:

1. **Hypericum Perforatum:** Indicated for nerve injuries with shooting or burning pains. Useful in cases of brachial neuritis or post-traumatic neuralgia. Helps in promoting nerve regeneration and pain relief.
2. **Causticum:** Beneficial for paralysis or weakness of upper limbs following nerve damage. Indicated when there is stiffness, contracture, or trembling.

Especially useful in long-standing nerve injuries.

3. **Rhus Toxicodendron**: Suitable for pain and stiffness in shoulder and arms, worse in cold and damp weather. Relieves muscular and ligamentous strain around the brachial plexus region. Often prescribed in inflammatory conditions with restlessness.
4. **Arnica Montana:** Indicated for traumatic injuries of nerves and soft tissues. Relieves soreness, bruised sensations, and post-surgical nerve pain. Supports faster recovery when taken early after trauma.
5. **Gelsemium Sempervirens:** Useful for weakness and heaviness in arms due to nerve involvement. Indicated in cases with trembling, dullness, and lack of coordination. Supports both functional and emotional aspects of nerve disorders. (7-8)

CONCLUSION:

The brachial plexus is a highly intricate nerve network that governs the motor and sensory functions of the upper limb. Its injury or disease can severely impair daily activities, leading to pain, weakness, or even paralysis. While surgical and rehabilitative measures remain primary management options, homeopathy can play a supportive role by alleviating pain, reducing inflammation, and enhancing the body's natural healing response. Remedies such as Hypericum, Causticum, Rhus tox, Arnica, and Gelsemium are frequently used in clinical practice to aid recovery. A holistic approach, combining modern medicine with homeopathy and physiotherapy, may provide better outcomes and improve quality of life for patients with brachial plexus disorders.

REFERENCES

1. Singh, Vishram. Textbook of Clinical Neuroanatomy, 5th Edition
2. Snell, R.S. Clinical Neuroanatomy, 7th Edition
3. Borley and Henry Gray, Gray's Anatomy: The Anatomical Basis of Clinical Practice.
4. Singh, Inderbir. Textbook Neuroanatomy, 2nd Edition
5. Sciencebasedmedicine.org/wp-content/uploads/2021/08/brachialplexuspalsy-e1628248807123.jpg
6. Dr Prakash, Erb's palsy and Klumpke's palsy - Etiology, Clinical features
7. Boericke W. New Manual of Homoeopathic Materia Medica and Repertory; 37th Edition.

How to Cite this Article- Singhal H., Brachial Plexus: Anatomy, Clinical Correlations, And Homeopathy. TUJ. Homo & Medi. Sci. 2025;8(3):69-73.