TANTIA UNIVERSITY JOURNAL OF HOMOEOPATHY AND MEDICAL SCIENCE
Website- www.tjhms.com Email : tujhms@tantiauniversity.com
Peer-Reviewed Journal, E-ISSN: 2581-8899, P-ISSN: 2581-978X
Volume 8|Issue 2|Apr. –June 2025|

REVIEW ARTICLE

THE ROLE OF HOMOEOPATHY IN OSTEOARTHRITIS: A REVIEW

Poonam Singh[1], Pranesh Kumar Singh[2]
[1]Sri Ganganagar Homoeopathic Medical College, Hospital and Research Institute, Sri Ganganagar, Rajasthan,
[2]State Ghazipur Homoeopathic Medical College & Hospital, Ghazipur

Received- 25/05/2025
Revised- 27/06/2025
Accepted– 30/06/2025

Key Word- OA, Cartilage degenartion, Osteophyte, Joint Disease, Homoeopathy.

Corresponding Author:- Poonam Singh, Professor, Sri Ganganagar Homoeopathic Medical College, Hospital and Research Institute, Sri Ganganagar, Rajasthan.

Abstract

Osteoarthritis (OA) is the most prevalent chronic joint disease, characterized by progressive cartilage degeneration, subchondral bone remodeling, osteophyte formation, and varying degrees of synovial inflammation. It represents a leading cause of pain and disability worldwide, especially among the elderly. This review summarizes current understanding of OA, including its epidemiology, risk factors, pathophysiology, clinical features, diagnostic approaches, and management strategies, with an emphasis on both conventional and complementary treatment options.

INTRODUCTION

Osteoarthritis (OA) is the most common form of arthritis — a chronic, degenerative joint condition where the protective cartilage that cushions the ends of your bones gradually wears down. Osteoarthritis is a degenerative joint disorder involving mechanical and biochemical factors that destabilize the normal coupling of cartilage synthesis and degradation. The disease primarily affects synovial joints and leads to progressive loss of function. Historically viewed as a "wear-and-tear" disease, OA is now recognized as a complex, whole-joint disorder influenced by inflammatory pathways, mechanical stress, and systemic metabolic changes.

Osteoarthritis (OA) is a degenerative joint disease marked by

progressive cartilage wear, subchondral bone changes, and varying degrees of inflammation. It is a leading cause of chronic pain and disability, particularly in older adults. Conventional medicine focuses on pain relief, improving function, and slowing disease progression through medications, physiotherapy, and surgery. However, many patients seek complementary approaches such as homeopathy to manage symptoms and improve quality of life.

OA commonly affects weight-bearing joints like the knees, hips, and spine, as well as smaller joints in the hands. Symptoms include joint pain, stiffness (especially after rest), swelling, and reduced mobility. Risk factors include age, obesity, joint injury, repetitive stress, and genetic predisposition.

Epidemiology

Global prevalence: OA affects over 500 million people worldwide, with knee OA being the most common form. Age distribution: Incidence increases significantly after the age of 50. Sex differences: Women are more commonly affected, especially post-menopause. Economic burden:OA is a major contributor to disability-adjusted life years (DALYs) and health care costs.

Risk Factors

- **Non-modifiable:** Age, genetics, sex, congenital joint abnormalities.
- **Modifiable:** Obesity, joint injury, occupational or sports-related overuse, muscle weakness, metabolic syndrome.

Pathophysiology-

OA is marked by an imbalance between anabolic and catabolic processes in cartilage and surrounding tissues.

- Cartilage damage: Cartilage normally provides a smooth, slippery surface for joint movement. In OA, it thins and breaks down. Loss of proteoglycans, collagen disruption, decreased chondrocyte function.
- Bone changes: The underlying bone may thicken, form bony spurs (osteophytes), or develop cysts.
- Subchondral bone: Increased bone turnover, sclerosis, cyst formation.
- Synovium: Mild to moderate inflammation with cytokine release (e.g., IL-1β, TNF-α).
- Osteophytes: Bony outgrowths formed as a compensatory response to instability.
- Inflammation: The joint lining (synovium) can become mildly inflamed, causing pain and stiffness.

CLINICAL FEATURES

- Joint pain: Worsens with activity, relieved by rest.
- Morning stiffness: Usually <30 minutes.

- Crepitus: Grinding sensation on movement.
- Decreased range of motion.
- Possible joint swelling and deformity in advanced cases.
- Pain (worse after activity or toward the end of the day)
- Joint stiffness (especially after rest, often called "gelling")

Commonly affected joints:

- Knees
- Hips
- Hands (finger joints, base of the thumb)
- Spine (neck or lower back)

DIAGNOSIS

- Clinical evaluation: Based on history and examination.
- Imaging:
  - X-ray: Joint space narrowing, osteophytes, subchondral sclerosis.
  - MRI: Early cartilage changes and bone marrow lesions.
  - Laboratory tests: Usually normal; used to exclude other arthritides.

MANAGEMENT

While there's no cure, treatment focuses on symptom relief, slowing progression, and maintaining joint function:

Non-pharmacological

- Lifestyle: Weight management, regular low-impact exercise (walking, swimming, cycling)
- Patient education and self-management programs.
- Exercise: Strength training, aerobic activity, flexibility exercises.
- Weight reduction to decrease joint load.
- Assistive devices: Braces, orthotics, canes.

Pharmacological

- Medications: Pain relievers (paracetamol), NSAIDs, topical agents
- First-line: Paracetamol (acetaminophen).
- Second-line: NSAIDs (oral or topical).
- Adjuncts: Duloxetine, topical capsaicin.
- Intra-articular Injections: Corticosteroids, hyaluronic acid.

Surgical

- Arthroscopy (limited role in OA).
- Osteotomy for joint realignment.
- Joint replacement (arthroplasty) in advanced disease.

Complementary and Alternative Therapies

- Nutritional supplements: Glucosamine, chondroitin (mixed evidence).

- Herbal remedies: Curcumin, Boswellia serrata.
- Homeopathy & acupuncture: Used by some patients; evidence remains limited but may help with symptom relief.

Homeopathy is a holistic system of medicine developed in the late 18th century by Dr. Samuel Hahnemann. It is based on the principle of "like cures like", where substances that produce symptoms in a healthy person are used in highly diluted forms to treat similar symptoms in illness. Remedies are chosen individually, based on the patient's physical symptoms, emotional state, and overall constitution.

In homeopathy, OA is not treated as a single uniform condition; instead, treatment is tailored to the individual's symptom pattern. The aim is to:

- Relieve pain and stiffness
- Improve joint mobility
- Reduce inflammation
- Support the body's natural healing capacity
- Address underlying tendencies toward degeneration

Commonly Used Homeopathic Medicines in Osteoarthritis

1. **Arnica Montana** – Soreness and bruised pain, as if beaten. Worse from touch, motion; better from lying down quietly. OA pain after overuse, trauma, or strain; joints feel tender to touch.For joint soreness and bruised pain, especially after overuse.

2. **Bryonia alba** –Pain aggravated by the slightest motion, relieved by complete rest. Worse in dry weather, from touch or pressure; better lying on the affected side. OA with hot, swollen joints and severe pain on movementFor joints that hurt with the slightest movement and feel better with rest.

3. **Rhus toxicodendron** - Stiffness and pain worse on first movement after rest ("rusty hinge"), improving with continued gentle motion. Worse in cold, damp weather; better with warmth and continued movement.OA with stiffness on waking, improved after moving around; often used in elderly patients with morning stiffness.

4. **Calcarea fluorica** – For bony outgrowths and joint deformities. Hard, bony outgrowths (osteophytes) and joint deformities. Worse in damp weather; better with warmth. OA with marked bone changes, stiffness, and reduced flexibility.

5. **Symphytum officinale** – Supports bone and cartilage health after injury. Promotes healing of bone and cartilage. OA following old fractures or joint injuries; helps repair damage to articular cartilage.

6. **Causticum-** Stiffness with contractures and weakness in tendons and ligaments. Worse in dry cold; better in damp weather and with heat. OA with deformities and restricted joint movement.

7. **Kali Carbonicum** - Sharp, stitching pains in joints, especially back and knees. Worse in cold weather, early morning; better with warmth. OA with swelling and back involvement, especially in elderly or those with a stooped posture.

8. **Ruta graveolens** - Deep, aching pain in tendons, ligaments, and periosteum. Worse from cold and damp; better from warmth. OA after repetitive strain or overuse of joints.

Integrating Homeopathy with Conventional Care

Homeopathy may be used alongside conventional OA management, especially for those seeking a gentler, holistic approach. It should not replace medical evaluation, particularly when joint damage is advanced or surgery is being considered. An integrated plan might include:

- Lifestyle changes (exercise, weight control)
- Physical therapy
- Nutritional support
- Carefully selected homeopathic remedies
- Medical monitoring for disease progression

CONCLUSION

Osteoarthritis is a multifactorial, progressive joint disease with significant individual and societal impact. While current treatments focus on symptom control and functional preservation, emerging therapies hold promise for altering disease progression. A multidisciplinary approach integrating patient education, lifestyle modification, pharmacotherapy, and in some cases surgery remains the cornerstone of OA management.

Homeopathy offers a patient-centered approach to osteoarthritis management, focusing on symptom relief and improving quality of life through individualized treatment. While scientific evidence is still evolving, many patients report subjective improvements in pain and mobility. Used alongside conventional therapies, homeopathy can be a supportive tool for those living with OA.

REFERENCES

1. Huang G, Zheng Y, Hong W, Qu X, Yang W, Cao H, Tian F, Lin H, Pei B, Chen B, Yang S, Qian D. Global, region and country burden of osteoarthritis at different sites in middle-aged and elderly populations from 1990 to 2021: a systematic analysis of the 2021 global burden of disease study. Front Med (Lausanne). 2025;12:1567303. doi:10.3389/fmed.2025.1567303
2. Xie X, Zhang K, Li Y, Li Y, Li X, Lin Y, et al. Global, regional, and national burden of osteoarthritis from 1990 to 2021 and projections to 2035: a cross-

sectional study for the Global Burden of Disease Study 2021. PLoS One. 2025;20(5)\:e0324296. doi:10.1371/journal.pone.0324296

3. Qiao L, Li M, Deng F, Wen X, Deng H, Xue Z, Xiang R, et al. Epidemiological trends of osteoarthritis at the global, regional, and national levels from 1990 to 2021 and projections to 2050 \[preprint]. medRxiv. 2024. doi:10.1101/2024.06.30.24309697
4. Scheuing W, Reginato A, Deeb M, Acer Kasman S. The burden of osteoarthritis: Is it a rising problem? *Best Pract Res Clin Rheumatol*. 2023;37:101836. doi:10.1016/j.berh.2023.101836
5. Chen X, Tang H, Lin J, Zeng R. Temporal trends in the disease burden of osteoarthritis from 1990 to 2019, and projections until 2030. *PLoS One*. 2023;18\:e0288561. doi:10.1371/journal.pone.0288561
6. Losina E, Weinstein A, Reichmann W, Burbine S, Solomon D, Daigle M, et al. Lifetime risk and age at diagnosis of symptomatic knee osteoarthritis in the US. *Arthritis Care Res (Hoboken)*. 2013;65:703–11. doi:10.1002/acr.21898
7. Litwic A, Edwards M, Dennison E, Cooper C. Epidemiology and burden of osteoarthritis. *Br Med Bull*. 2013;105:185–99. doi:10.1093/bmb/lds038

How to Cite this Article- Singh P., Singh P. K., Chauhan G., The Role Of Homoeopathy In Osteoarthritis: A Review. TUJ. Homo & Medi. Sci. 2025;8(2):60-65.

Conflict of Interest: None **Source of Support: Nil**